

# Breathe Easy With Your 4 Puffagins a Day

Write in the times of day that 4 Puffagins might fit in your routine.

\_\_\_\_ : \_\_\_\_ AM / PM

## YOUR 4 PUFFAGINS WORK TOGETHER TO HELP YOU REMEMBER YOUR PUFFS

The usual starting dose for ATROVENT HFA is 2 puffs, 4 times a day. And it's a good idea to spread these doses throughout your day. But you should not take more than 6 doses (12 puffs) in 24 hours.

### INDICATION FOR USE

ATROVENT HFA (ipratropium bromide HFA) is a prescription maintenance treatment for bronchospasm (airway narrowing) in patients with chronic obstructive pulmonary disease (COPD), including chronic bronchitis and emphysema.

### IMPORTANT SAFETY INFORMATION

Do not use ATROVENT HFA (ipratropium bromide HFA) if you are allergic to ipratropium or any of the ingredients in ATROVENT HFA or to atropine or similar drugs.

ATROVENT HFA is not a rescue medicine and should not be used for treating sudden breathing problems.

Please see complete Important Safety Information on back and read accompanying full Prescribing Information, including Instructions for Use.

**ATROVENT** HFA  
(ipratropium bromide HFA)  
INHALATION AEROSOL

| the  
Puffagins

## IMPORTANT SAFETY INFORMATION

Do not use ATROVENT HFA (ipratropium bromide HFA) if you are allergic to ipratropium or any of the ingredients in ATROVENT HFA or to atropine or similar drugs.

ATROVENT HFA is not a rescue medicine and should not be used for treating sudden breathing problems. Your doctor may give you other medicine to use for sudden breathing problems.

Allergic reactions may occur, including itching, swelling of the face, lips, tongue, or throat (involving difficulty in breathing or swallowing), rash, hives, bronchospasm (airway narrowing), or anaphylaxis. Some of these may be serious. If you experience any of these symptoms, stop taking ATROVENT HFA at once and call your doctor or get emergency help.

ATROVENT HFA can cause the narrowing of the airways to get worse (paradoxical bronchospasm) which may be life threatening. If this happens, stop taking ATROVENT HFA at once and call your doctor or get emergency help.

ATROVENT HFA may increase eye pressure which may cause or worsen some types of glaucoma.

Do not get the spray into your eyes. The spray may cause eye pain or discomfort, blurred vision, enlarged pupils, seeing halos or colored images along with red eyes. If you have any of these symptoms, stop taking ATROVENT HFA and call your doctor right away.

Dizziness and blurred vision may occur with ATROVENT HFA. Should you experience these symptoms, use caution when engaging in activities such as driving a car or operating appliances or machinery.

ATROVENT HFA may cause difficulty with urination. Symptoms may include difficulty passing urine and/or painful urination. If you have any of these symptoms, stop taking ATROVENT HFA and call your doctor right away.

The most common side effects reported with use of ATROVENT HFA were bronchitis, COPD flare-up (exacerbation), shortness of breath and headache.

Tell your doctor about all medicines you are taking, including eye drops. Ask your doctor if you are taking any anticholinergic medicines because taking them together with ATROVENT HFA can increase side effects.

## INDICATION FOR USE

ATROVENT HFA (ipratropium bromide HFA) is a prescription maintenance treatment for bronchospasm (airway narrowing) in patients with chronic obstructive pulmonary disease (COPD), including chronic bronchitis and emphysema.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.FDA.gov/medwatch](http://www.FDA.gov/medwatch) or call 1-800-FDA-1088.

**Please see accompanying full Prescribing Information, including Instructions for Use.**

**ATROVENT<sup>®</sup> HFA**  
(ipratropium bromide HFA)  
INHALATION AEROSOL



Copyright ©2017 Boehringer Ingelheim Pharmaceuticals, Inc.  
All rights reserved. (05/17) PC-04194